

L1014771 IREL/10T PPE4058713



**Klerat® Waxblocks**

BRAND LOGO/NAME  
LOGO/NOM DE MARQUE  
LOGO/NOMBRE DEL PRODUCTO

**Rodenticide**

**syngenta®**

Contains 0.05 g/kg (0.005% w/w) brodifacoum

**Klerat® Waxblocks is a ready-for-use waxblock bait containing 0.05 g/kg brodifacoum (0.005 % w/w), for use as a rodenticide for the control of rats and mice indoors, outdoors around buildings and in sewers by professionals for the protection of public health, animal health, stored products and materials.**

FRONT PAGE

TEXT AREA  
214 x 115 mm

Product names marked ® or ™, the ALLIANCE FRAME, the SYNGENTA Logo and the PURPOSE ICON are Trademarks of a Syngenta Group Company

**10 KG e**

Batch number: Refer to packaging  
Expiry date: Refer to packaging

7

DESCRIPTION:  
BOOKLET 4-16 PG 275 X 122 MM

SCALE:  
1:1

ISSUE NO:  
SYN002

ISSUE DATE:  
24.10.2014

**4032233**



Rodenticide

**DIRECTIONS FOR USE:**

For use as a rodenticide only. Klerat® Waxblocks are active against black rats, brown rats and mice. Always read the label before use and follow the instructions provided. Avoid all unnecessary exposure; in particular avoid ingestion. Product must be handled in a safe manner. Do not decant product into unlabelled containers. A thorough survey of the infested area is essential, particularly in secluded and sheltered places, to determine the extent of the infestation. The product may be applied by professional users only when appropriate chemical resistant gloves (EN 374) are used.

**Resistance Information:**

The resistance status of the target population should be taken into account when considering the choice of rodenticide to be used. In those areas where evidence of resistance to specific active ingredients is suspected, avoid their use. Where resistance to brodifacoum has been shown or suspected, resistance management strategies should be employed. To control the spreading of resistance, it is advisable to alternate baits containing different anticoagulant active ingredients.

**Use indoors and outdoors (around buildings):**

**Rats:** Secure 20-60g of bait in covered tamper resistant bait stations or covered bait points spaced 10m apart (5m apart in high infestation areas) in areas where rats are active.

**Mice:** Secure 20g of bait in covered tamper resistant bait stations or covered bait points spaced 5m apart (2m apart in high infestation areas) in areas where mice are active.

**Use in sewers:**

**Rats:** Secure 20-200g of bait per station to available structures to ensure blocks are not washed away. Initial inspection of bait points after 3 or 4 days is recommended. Replace any bait that has been consumed. Repeat approximately 4 days later and then weekly. Replace contaminated or spoiled bait until consumption has stopped. Repeat treatment in situations where there is evidence of new infestation (e.g. fresh tracks or droppings). If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait points. Do not increase the bait point size. The product must never be placed indiscriminately. Bait must be securely deposited in baiting stations or other coverings so as to minimise the risk of consumption by other animals or children. Where possible, secure baits so that they cannot be dragged away. When used in sewers, bait should be placed so it is protected from flooding or displacement. Prevent access to bait by children, birds and non-target animals (particularly dogs, cats, pigs and poultry). Use bait containers clearly marked "poison" at all surface baiting points. It is illegal to use this product for the intentional poisoning of non-target, beneficial and protected animals. Wash hands and face after application and use of the product, and before eating, drinking or smoking. For professional users the use of appropriate personal protective equipment (PPE) is advised. Search for and remove dead rodents at frequent intervals during treatment (unless used in sewers), at least as often as when baits are checked and/or replenished. Daily inspection may be required in some circumstances. Dispose of dead rodents in accordance with local requirements. Poisoned rodents may be disposed of by the waste producer at an incinerator or landfill permitted to accept that type of waste, or collected by a registered waste carrier and taken for disposal at a suitably permitted site.

Revuez s'il vous plait avant l'impression. Por favor, eliminar antes de imprimir.

**Use in public areas:** When the product is being used in public areas and tamper resistant bait stations are not used, the areas treated must be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anti-coagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits. When tamper resistant bait stations are used, they should be clearly marked to show that they contain rodenticides and that they should not be disturbed.

**Use in sewers:** For use in sewers where there is no risk to children, companion animals and non-target species blocks should be secured to available structures by wire to ensure the block is not washed away. Unless under the supervision of a pest control operator or other competent person, do not use anticoagulant rodenticides as permanent baits. In most cases, anticoagulant bait should have achieved control within 35 days. Should activity continue beyond this time, the likely cause should be determined.

**Precautions:**

- Keep out of reach of children.
- Read label before use.
- Keep/Store away from food, drink and animal feedingstuffs.
- Do not get on skin.
- Do not eat, drink or smoke when using this product.
- Avoid release to the environment.
- Wear protective gloves.
- IF SWALLOWED: Immediately call a POISON CENTRE/doctor.
- Store locked up in a closed container.
- Dispose of contents/container in accordance with national regulations.
- Wash hands and exposed skin before meals and after use.

THESE PAGES CAN BE ADDED OR REMOVED AS NECESSARY TO CREATE A 4, 8, 12, OR 16 PAGE BOOKLET (INCLUDING GLUE PAGE)

TEXT AREA  
214 x 115 mm

THE PENULTIMATE PAGE IS PERMANENTLY AFFIXED TO THE BOTTLE, REPEAT LEGAL REQUIREMENTS IF NECESSARY (eg. FRONT PAGE)

IF USING A REPEAT PAGE, PLEASE SWITCH ON VISIBILITY OF 'REPEAT PAGE' LAYER IN INDESIGN FILE

**Storage:**

Store in a cool, dry, well-ventilated place. Store away from light. Keep only in original container. A shelf life of 2 years is supported.

**Disposal:**

Remove all baits after treatment and dispose of them in accordance with local requirements. Dead rodents and bait found outside bait stations should be disposed of using protective gloves. This material and its container must be disposed of safely. Dispose of packaging, remains of unused product and dead rodents in accordance with local regulations. If required, consult a professional waste operator or local authority.

Harmful to wildlife.

To avoid risks to human health and the environment, comply with the instructions for use.

Use Biocides Safely and Sustainably.

It is illegal to use this product for uses or in a manner other than that prescribed on the label. NOT for Amateur Sale.

**First Aid:**

In case of accident, or if you feel unwell, seek medical advice immediately (show the label where possible). In case of contact with skin, wash with soap and water. Remove and launder any contaminated clothing. In case of contact with eyes, remove contact lenses if present and rinse the eye slowly and gently with water for 15-20 minutes. Seek medical advice immediately. In case of ingestion or if swallowed seek medical advice immediately.

TEXT AREA  
214 x 115 mm

THE PENULTIMATE PAGE IS PERMANENTLY AFFIXED TO THE BOTTLE, REPEAT LEGAL REQUIREMENTS IF NECESSARY (eg. FRONT PAGE)

IF USING A REPEAT PAGE, PLEASE SWITCH ON VISIBILITY OF 'REPEAT PAGE' LAYER IN INDESIGN FILE

**Note to Doctor/Physician:**

Antidote vitamin K1 (under medical supervision).  
Effects result from the inhibition of blood coagulation, through interaction with vitamin K1 recycling. Poisoning symptoms may include: Bruising easily, nose or gum bleeds, blood in stools or urine, excessive bleeding from minor cuts or abrasions. Note that poisoning symptoms may develop over the course of several days.  
Brodifacoum is an anticoagulant which may produce bleeding; this may be delayed for several days after exposure. If there is no active bleeding the INR (prothrombin time) should be measured on presentation and 48-72 hours after exposure. If the INR is greater than 4, administer Vitamin K1 (phytomenadione) 5-10 mg by slow intravenous injection (100 µg/kg body weight for a child). Treatment with phytomenadione (orally or intravenously) may be required for several weeks.

**Poison Centre information and reporting of incidents:**

Poison Information: For information or to report a poisoning incident contact;  
The National Poisons Information Centre, Beaumont Hospital, Dublin (01-8092166).  
Retain the label for reference.

**Authorisation Holder and Marketing Company:**

Syngenta Crop Protection AG, Schwarzwaldallee 215, CH-4058 Basel, Switzerland  
Emergency Telephone: +44(0)1484 538444  
Authorisation number: IE/BPA 70295

**IF USING A REPEAT PAGE, PLEASE SWITCH ON VISIBILITY  
OF 'REPEAT PAGE' LAYER IN INDESIGN FILE**

L1014771 IREL/10T PPE4058713

**Klerat® Waxblocks**  
Rodenticide  
(INCLUDING GLUE PAGE)

**syngenta®**

Contains 0.05 g/kg (0.005% w/w) brodifacoum

**Klerat® Waxblocks is a ready-for-use waxblock bait containing 0.05 g/kg brodifacoum (0.005 % w/w), for use as a rodenticide for the control of rats and mice indoors, outdoors around buildings and in sewers by professionals for the protection of public health, animal health, stored products and materials.**

Product names marked ® or ™, the ALLIANCE FRAME, the SYNGENTA Logo and the PURPOSE ICON are Trademarks of a Syngenta Group Company

**10 KG e**

**IF USING A REPEAT PAGE, PLEASE SWITCH ON VISIBILITY  
OF 'REPEAT PAGE' LAYER IN INDESIGN FILE**

Batch number: Refer to packaging  
Expiry date: Refer to packaging

NO TEXT HERE

GLUE PAGE

PAGE NUMBER SEQUENCING IS DETERMINED BY THE FINAL LEADING EDGE.  
RIGHT HAND STARTS FROM BACK OF BOOKLET. LEFT HAND STARTS FROM FRONT OF BOOKLET.  
FIRST (FRONT) PAGE IS ALWAYS FIXED ON TOP OF BOOKLET.