



Pybuthrin[®] 33

INSECTICIDE

- ✓ Ideal for hot / thermal fogging
- ✓ Natural active ingredient
- ✓ Biodegradable and no residual activity

Pybuthrin[®] 33

Contains: Pyrethrins including cinerins 0.38% w/w (3 g/l), odourless kerosine and synergised with piperonyl butoxide.

DANGER



May be fatal if swallowed and enters airways.
Causes skin irritation.

Very toxic to aquatic life with long lasting effects.

Wear protective gloves/protective clothing/eye protection/face protection.

If exposed or concerned: Call a POISON CENTER or doctor/physician.


Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

Contains pyrethrins including cinerins.

May produce an allergic reaction.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS 93411, HSE 5106



5 L e

STATUTORY CONDITIONS RELATING TO USE.

FOR USE ONLY AS AN INSECTICIDE.

FOR PROFESSIONAL USE in Public Hygiene/Nuisance.

Dose :

Space spray:

Indoors: 1 litre of product per 3000 m³.

Outdoors: Up to 25 litres per ha / 1 litre per 2000 m².

Surface (contact) spray: 5 litres of product per 100 m².

The (COSHH) Control of Substances Hazardous to Health Regulations 2002 may apply to the use of this product at work.

Engineering control of operator exposure must be used where reasonably practicable in addition to the following items of personal protective equipment.
Wear suitable protective clothing (COVERALLS) SYNTHETIC RUBBER/PVC GLOVES AND EYE PROTECTION when using.

DO NOT BREATHE FOG OR SPRAY MIST. Otherwise wear respiratory protective equipment and eye protection (see HSE Guidance Booklet HS (G) 53: "The Selection, Use and Maintenance of Respiratory Protective Equipment – A Practical Guide").

AVOID EXCESSIVE CONTAMINATION OF COVERALLS AND LAUNDRER REGULARLY.

However engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

If swallowed, do not induce vomiting: seek medical advice immediately and show this container or label.

DO NOT CONTAMINATE FOODSTUFFS, EATING UTENSILS OR FOOD CONTACT SURFACES.

COVER WATER STORAGE TANKS before application.

WHEN USING DO NOT EAT DRINK OR SMOKE.

Keep away from sources of ignition - No smoking

Do not apply in the presence of naked flames, hot surfaces or unprotected electrical equipment.

UNPROTECTED PERSONS AND ANIMALS SHOULD BE KEPT AWAY FROM TREATED AREAS UNTIL SURFACES ARE DRY.

Do not contaminate ground, waterbodies or watercourses with chemicals or used container.

HARMFUL TO FISH AND OTHER AQUATIC LIFE REMOVE OR COVER ALL FISH TANKS AND BOWLS before application.

HAZARDOUS TO BEES.

This material and its container must be disposed of in a safe way.

AVOID ALL CONTACT WITH PLANT LIFE.

All bats are protected under the Wildlife and Countryside Act (1981) or the Wildlife (Northern Ireland) Order (1985). Before treating any structure used by bats, consult English Nature, Scottish Natural Heritage, The Countryside Council for Wales or Department of the Environment for Northern Ireland.

FOR IRELAND: ALL BATS ARE PROTECTED UNDER SCHEDULE 5 OF THE WILDLIFE ACTS (1976 AND 2000). BEFORE TREATING ANY STRUCTURE USED BY BATS, CONSULT THE NATIONAL PARKS AND WILDLIFE SERVICE.

READ ALL PRECAUTIONS BEFORE USE

PCS 93411, HSE 5106

When stored in a dry place at temperature not exceeding 25°C Pybuthrin 33 has a shelf life of 2 years.
This product is approved under The Control of Pesticides Regulations (1986) for use as directed (HSE 5106).

A pyrethrin spray combining rapid knockdown and kill with non-persistence making it suitable for use in food handling premises. Pybuthrin 33 will also flush out insects from their harboursages.

For the control of:

Flying Insects e.g. flies, wasps (and wasps nests), and stored product moths and mosquitoes.

Crawling Insects e.g. cockroaches and stored product insects.

DIRECTIONS FOR USE

A ready-to-use space spray/fogger and surface (contact spray) for use in the following situations.

Domestic Situations: Private housing.

Food Handling:

Processing: food manufacture, kitchens, slaughterhouses etc.

Storage: food retailers, warehouses, raw material stores etc.

Preparation: restaurants, commercial kitchens etc.

Public Buildings:

Hospitals (excluding occupied wards), hotels, public baths, municipal buildings, churches, halls, community centres, cinemas etc.

Outdoors

Refuse tips, sewage works (excluding filter beds), amenity areas etc.

RATES OF APPLICATION

Where	Spraying Equipment	Volume	Area	Other Information
Space Spray Indoor	Apply with any suitable hand, mechanically or electrically operated sprayer capable of producing a fine mist or fog.	1 litre	3000 m ³	Close windows and doors before spraying and for at least 30 minutes afterwards. Distribute evenly as a fine mist or fog. For moth infestations, treatment at weekly intervals is recommended until infestations are brought under control. More frequent applications may be necessary for heavy infestations. Spray as frequently as the situation demands for flying insect control.
Space Spray Outdoor (Urban/Industrial sites)		1 litre 1 litre	2000 m ² for large area 400 m ² for small area	The best conditions for outdoor spraying occur in the early morning or late afternoon and evening when conditions are most likely to be still. Spray as frequently as the situation demands.
Surface (contact) Spray	Apply with any suitable hand, mechanically or electrically operated sprayer capable of producing a coarse spray. Use only equipment adjusted to give a coarse spray.	1 litre	20 m ²	Sufficiently dampen the insects. Apply directly to the insects present on the fabric of warehouses, granaries, mills, bakeries and similar situations, or insects on the surfaces of bags and cases containing stored food products. Special attention should be paid to cracks and crevices and other insect harboursages.
Wasp Nests		Apply liberally and directly to nest or to entrance holes of concealed nests	Wasp Nest directly or entrance holes of concealed nests	Apply at dusk or when activity around the nest is quiet.

SPRAYING EQUIPMENT

As a space spray:

Apply with any suitable hand, mechanically or electrically operated sprayer capable of producing a fine mist or fog.

As a surface (contact) spray:

Apply with any suitable hand, mechanically or electrically operated sprayer capable of producing a coarse spray. Use only equipment adjusted to give a coarse spray.

WARNINGS

Avoid inhalation of spray and contact with skin and eyes.

Do not spray the moving parts of any machinery, electric motors and switchgear.

Do not spray and avoid spilling on plastics, synthetic tiles or light coloured surfaces.

This spray will not harm ordinary fabrics or affect commonly used dyes, but before spraying delicate or expensive materials test first by rubbing a little spray on some inconspicuous spot.

Polished surfaces may lose their gloss until re-polished.

This spray is NOT suitable for use on plants.

It is illegal to disturb bats or their roosts and this may influence the use of Pybuthrin 33 in roof spaces etc.

No treatment should be made within a roof space without first ascertaining whether bats use the site as a roost.

PRECAUTIONS

ENSURE ADEQUATE VENTILATION WHEN APPLYING.

VENTILATE TREATED AREAS THOROUGHLY after application.

WASH HANDS AND EXPOSED SKIN before meals and after use.

WASH ANY SPLASHES/CONTAMINATION from skin or eyes immediately.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP OUT OF THE REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed in a safe place.

Use appropriate containment to avoid environmental contamination.

RA3 - Pybuthrin 33 - GB - 5 L - GB05820307H - ARTICLE 05869284

Approval Holder:
Bayer CropScience Limited

WHO-classification:
U (Unlikely to present acute hazard in normal use).

For 24 hour emergency information
contact Bayer CropScience Ltd.
From the UK: 0800 220876
From the Republic of Ireland: Freephone: 1800 409399

Pybuthrin® is a registered trademark of Bayer.
© Copyright of Bayer 2003 – All rights reserved.

Bayer

Authorisation holder:
Bayer CropScience Limited
230 Cambridge Science Park
Milton Road, Cambridge CB4 0WB
From the UK and the Republic of Ireland:
08000 1214 9451
www.pestcontrol-expert.com
or MSDS and larger label