

sakarati[®] D LIQUID BAIT

0.005%W/W (0.05g/kg) Difenacoum

Professional use only in and around buildings



APPLICATION RATE (S) AND FREQUENCY:

MICE: Up to 100ml bait in bait station every 2-5 metres
RATS: Up to 250ml bait in bait station every 5-10 metres

The dispenser must always be placed in a correctly labelled, tamper-resistant bait station.

Bait stations are manually placed in the rodent infested area. Bait station should be fixed to the ground.

The product must never be placed indiscriminately

PROFESSIONAL USE:

To be used only by professional users holding certification demonstrating compliance with UK rodenticide stewardship regime requirements.

Read the label before use. Using this product in a manner that is inconsistent with the label may be an offence. Refer to the CRRU UK Code of Best Practice (or equivalent) for guidance.

When this product is supplied to a user for the control of rodents, it shall only be supplied to a professional user holding certification demonstrating compliance with UK rodenticide stewardship regime requirements.

The dispenser must always be placed in a correctly labelled, tamper-resistant bait station.

Bait stations must be securely deposited in a way so as to minimise the risk of consumption by other animals or children.

Bait bottles will be opened by removing the childproof-cap without breaking the membrane, the roll-on dispenser will then be connected to the bottle. Once inverted, the bait bottle and connected dispenser will be placed in the bait station.

The bait station will be fixed to the ground and the treatment signposted.

Do not use in areas where food/feed, cooking utensils or food processing surfaces could come into contact with the product, or be contaminated by it.

Do not eat, drink or smoke during the handling of the product and wash hands after use.

In order to prevent resistance and secondary poisoning of non-target animals, do not use the product as permanent bait.

To minimise problems of resistance and the risk of primary poisoning the product should not be used as permanent bait to prevent the appearance of rodents.

The treatment has to be alternated with other kinds of active substance in order to avoid resistance and because of cross resistance occurrence to rodenticides.

Do not clean the bait stations with water between applications.

In case of accidental spillage of the liquid, dispose of the bait station as hazardous waste.

Do not throw the product on the ground.

Do not place the product in water courses or sinks, do not pour down the drain.

Do not open the bottle, unless attaching to the roll on dispenser.

Collect dead rodents and unused baits during all control operations to minimise the risk of secondary poisoning.

Hazardous for wildlife.

Wear protective chemical resistant gloves during product handling phase (nitrile to EN374)

Use as part of an integrated pest management strategy (to include measures of chemical, physical and hygienic control).

The level of efficacy should be monitored with periodic checking, cases of reduced efficacy should be investigated for evidence of resistance.

In order to minimise the risk of secondary poisoning (consumption by companion animals and other non-target animals), search for and remove dead rodents at frequent intervals during treatment, as least as often as when baits are checked and/or replenished.

PARTICULARS OF LIKELY DIRECT OR INDIRECT EFFECTS, FIRST AID INSTRUCTIONS AND EMERGENCY MEASURES TO PROTECT THE ENVIRONMENT:

Basic vital support measures

Move the person away from the contaminated area and remove any contaminated or spattered clothing.

If in eyes: rinse cautiously with water for 15 minutes. Remove contact lenses if present.

If on skin: Wash with plenty of soap and water, without rubbing.

If swallowed: Do not induce vomiting, unless told to do so by poison centre or a health care professional.

Keep the patient at rest and maintain body temperature.

Check breathing. If necessary, give artificial respiration.

If necessary take the person to a hospital and show the label or packaging when possible.

DO NOT LEAVE POISONED PERSON ALONE UNDER ANY CIRCUMSTANCE

Clinical symptoms

Bleeding diathesis, derived from antiproteolytic effect- prolonged prothrombin time- that may become evident at minimum 24-maximum 72

hours (A prothrombin time when the patient is admitted in the hospital, does not exclude the diagnosis)

Medical advice for doctors and medical staff

If less than 2 hours has passed from the intake, gastric emptying must be performed, and activated carbon dispensed (25g)

Antidote: Vitamin K1 (phytonadione)

Check the prothrombin time

Treat symptomatically

IF MEDICAL ADVICE IS NEEDED, HAVE PRODUCT CONTAINER OR LABEL AT HAND AND CONTACT THE POISON CENTRE

To report human poisoning incidents call the relevant national poison information centre. Include information on the product authorisation number, product trade name and active substance. Where possible provide a copy of the label or safety data sheet.

INSTRUCTIONS FOR SAFE DISPOSAL OF THE PRODUCT AND ITS PACKAGING:

Remove all baits after treatment and dispose of them in accordance with local requirements.

Dead rodents and bait found outside bait stations should be disposed of using protective gloves.

Professionals: In the UK for information on disposal contact the Environment Agency (<http://www.environment-agency.gov.uk>) or SEPA (<http://www.sepa.org.uk>).

CONDITIONS OF STORAGE AND SHELF-LIFE OF THE PRODUCT UNDER NORMAL CONDITIONS OF STORAGE:

Keep in a cool, dry, well-ventilated area.

Store only in original container.

A shelf life of up to 2 years is supported

Store away from light.

Store out of reach of children and companion animals.

AUTHORISATION NO-UK-2017-1113

AUTHORISATION HOLDER:

Killgerm Chemicals Limited
Wakefield Road
Ossett
West Yorkshire
WF5 9AR

Date of authorisation; 31st October 2017

Expiry date of authorisation; 31st August 2020

TARGET ORGANISMS:
Brown Rat (*Rattus norvegicus*)
House Mouse (*Mus Musculus*)



DANGER

MAY DAMAGE FERTILITY OR THE UNBORN CHILD
MAY CAUSE DAMAGE TO ORGANS THROUGH PROLONGED OR REPEATED EXPOSURE

PRECAUTIONARY STATEMENTS:

KEEP OUT OF REACH OF CHILDREN.

READ LABEL BEFORE USE.

IF MEDICAL ADVICE IS NEEDED, HAVE PRODUCT CONTAINER OR LABEL AT HAND.

STORE LOCKED UP.

DO NOT EAT, DRINK OR SMOKE WHEN USING THIS PRODUCT.

WEAR PROTECTIVE GLOVES.

IF SWALLOWED: IMMEDIATELY CALL A POISON CENTRE OR DOCTOR/PHYSICIAN.

Contents:
50 x 100ml e
www.killgerm.com