

# Dismate PE

Management by confusion!

For professional use only.  
For indoor use only.  
For use only as an attractant.  
To avoid risk to man and the environment,  
comply with the instructions for use.

The COSHH (Control of Substances  
Hazardous to Health) Regulations 2002  
(as amended) apply to the use of this  
product at work.

DO NOT CONTAMINATE FOODSTUFFS,  
EATING UTENSILS OR FOOD CONTACT  
SURFACES.

Wash hands and exposed skin  
thoroughly after handling.

 **UK-2016-1028**  
 **EU-0024354-0000**

**UK authorisation holder:** Russell IPM,  
Unit 45, First Avenue, Deeside Industrial park,  
Deeside, Flintshire, UK CH5 2NU.

**EU authorisation holder:** Russell IPM Europe  
limited, Bond house 9/10 Bridge street,  
Lower Dublin 8, Dublin , Ireland.

**Active substance:**  
1433 – (9Z,12E)-tetradeca-9,  
12-dien-1-yl acetate 0,837%

**Type of Formulation:**  
VP-Vapour releasing product

## FIRST AID

**IF IN EYES:** If symptoms occur rinse with  
water. Remove contact lenses, if present  
and easy to do. Call a **POISON CENTRE**  
or a doctor.

**IF INHALED:** If symptoms occur call a  
**POISON CENTRE** or a doctor.

**IF ON SKIN:** Wash skin in water.  
If symptoms occur call a **POISON CENTRE**  
or a doctor.

**IF SWALLOWED:** If symptoms occur call  
a **POISON CENTRE** or a doctor.

For professional use only.  
For indoor use only.  
For use only as an attractant.  
To avoid risk to man and the environment,  
comply with the instructions for use.


The COSHH (Control of Substances  
Hazardous to Health) Regulations 2002  
(as amended) apply to the use of this  
product at work.

DO NOT CONTAMINATE FOODSTUFFS,  
EATING UTENSILS OR FOOD CONTACT  
SURFACES.

Wash hands and exposed skin  
thoroughly after handling.

# Dismate PE

Management by confusion!

 **UK-2016-1028**  
 **EU-0024354-0000**

**UK authorisation holder:** Russell IPM,  
Unit 45, First Avenue, Deeside Industrial park,  
Deeside, Flintshire, UK CH5 2NU.

**EU authorisation holder:** Russell IPM Europe  
limited, Bond house 9/10 Bridge street,  
Lower Dublin 8, Dublin , Ireland.

**Active substance:**  
1433 – (9Z,12E)-tetradeca-9,  
12-dien-1-yl acetate 0,837%

**Type of Formulation:**  
VP-Vapour releasing product

## FIRST AID

**IF IN EYES:** If symptoms occur rinse with  
water. Remove contact lenses, if present  
and easy to do. Call a **POISON CENTRE**  
or a doctor.

**IF INHALED:** If symptoms occur call a  
**POISON CENTRE** or a doctor.

**IF ON SKIN:** Wash skin in water.  
If symptoms occur call a **POISON CENTRE**  
or a doctor.

**IF SWALLOWED:** If symptoms occur call  
a **POISON CENTRE** or a doctor.

**Target insects:**  
Indian meal moths (*Plodia interpunctella*)  
Mediterranean flour moths (*Ephesia*  
*Khueniella*)

**Where to use:**  
Cereal processing, bakeries, confectionary  
manufacturing facilities and dried food  
processing/storage facilities.

**Application:**  
One dispenser every 8m x 8m at 2m height  
(400-500 cubic meter). Change dispensers  
every three months.

**Target insects:**  
Indian meal moths (*Plodia interpunctella*)  
Mediterranean flour moths (*Ephesia*  
*Khueniella*)

**Where to use:**  
Cereal processing, bakeries, confectionary  
manufacturing facilities and dried food  
processing/storage facilities.

**Application:**  
One dispenser every 8m x 8m at 2m height  
(400-500 cubic meter). Change dispensers  
every three months.

Approved for the use in the presence of exposed (unpacked) food.

Approved for the use in the presence of exposed (unpacked) food.

**Dismate PE** acts by disrupting the mating  
between male and female moths. The  
populations of the pest insects are  
reduced as adult insects die and are not  
replaced by the next generation.

**Dismate PE** is a vapourising dispenser in  
ready to use pouch. Keep the dispensers  
sealed in outer foil bag packaging until use.

**Store at temperatures not exceeding 4°C.**  
**Shelf life of up to 2 years.**

**Pack quantity: 5 or 100**

**DIRECTIONS FOR USE:**  
Plot out a layout for the approximate  
placement of the dispensers before deploying  
them. The dispensers should be placed one  
unit in every 400-500m<sup>3</sup>. Where obstacles or  
obstructions are present the dispensers can  
be placed to accommodate the layout and air  
dynamics of the factory.  
The dispensers should be positioned 2-3  
metres above floor and fastened securely  
using the supplied wall mounting bracket.  
Monitoring traps are recommended before,  
during and after treatment to assess

adult moth populations as part of a pest  
management program.  
Dispensers should be replaced every 3 months.  
Dispose of contents/container to national  
Regulations.  
This product has a slow mode of action and  
is intended for long-term use. The product  
doesn't have an immediate effect on the  
population of the food moths (it may take  
3-12 months to notice any significant  
reduction in moth population).  
Control of food moths is achieved over a period  
of time with continuous use of the product.

**Dismate PE** acts by disrupting the mating  
between male and female moths. The  
populations of the pest insects are  
reduced as adult insects die and are not  
replaced by the next generation.  
**Dismate PE** is a vapourising dispenser in  
ready to use pouch. Keep the dispensers  
sealed in outer foil bag packaging until use.

**Store at temperatures not exceeding 4°C.**  
**Shelf life of up to 2 years.**

**Pack quantity: 5 or 100**

**DIRECTIONS FOR USE:**  
Plot out a layout for the approximate  
placement of the dispensers before deploying  
them. The dispensers should be placed one  
unit in every 400-500m<sup>3</sup>. Where obstacles or  
obstructions are present the dispensers can  
be placed to accommodate the layout and air  
dynamics of the factory.  
The dispensers should be positioned 2-3  
metres above floor and fastened securely  
using the supplied wall mounting bracket.  
Monitoring traps are recommended before,  
during and after treatment to assess

adult moth populations as part of a pest  
management program.  
Dispensers should be replaced every 3 months.  
Dispose of contents/container to national  
Regulations.  
This product has a slow mode of action and  
is intended for long-term use. The product  
doesn't have an immediate effect on the  
population of the food moths (it may take  
3-12 months to notice any significant  
reduction in moth population).  
Control of food moths is achieved over a period  
of time with continuous use of the product.

# Dismate PE

Management by confusion!

For professional use only.  
For indoor use only.  
For use only as an attractant.  
To avoid risk to man and the environment,  
comply with the instructions for use.

The COSHH (Control of Substances  
Hazardous to Health) Regulations 2002  
(as amended) apply to the use of this  
product at work.

DO NOT CONTAMINATE FOODSTUFFS,  
EATING UTENSILS OR FOOD CONTACT  
SURFACES.

Wash hands and exposed skin  
thoroughly after handling.

 **UK-2016-1028**  
 **EU-0024354-0000**

**UK authorisation holder:** Russell IPM,  
Unit 45, First Avenue, Deeside Industrial park,  
Deeside, Flintshire, UK CH5 2NU.

**EU authorisation holder:** Russell IPM Europe  
limited, Bond house 9/10 Bridge street,  
Lower Dublin 8, Dublin , Ireland.

**Active substance:**  
1433 – (9Z,12E)-tetradeca-9,  
12-dien-1-yl acetate 0,837%

**Type of Formulation:**  
VP-Vapour releasing product

## FIRST AID

**IF IN EYES:** If symptoms occur rinse with  
water. Remove contact lenses, if present  
and easy to do. Call a **POISON CENTRE**  
or a doctor.

**IF INHALED:** If symptoms occur call a  
**POISON CENTRE** or a doctor.

**IF ON SKIN:** Wash skin in water.  
If symptoms occur call a **POISON CENTRE**  
or a doctor.

**IF SWALLOWED:** If symptoms occur call  
a **POISON CENTRE** or a doctor.

For professional use only.  
For indoor use only.  
For use only as an attractant.  
To avoid risk to man and the environment,  
comply with the instructions for use.



The COSHH (Control of Substances  
Hazardous to Health) Regulations 2002  
(as amended) apply to the use of this  
product at work.

DO NOT CONTAMINATE FOODSTUFFS,  
EATING UTENSILS OR FOOD CONTACT  
SURFACES.

Wash hands and exposed skin  
thoroughly after handling.

# Dismate PE

Management by confusion!

 **UK-2016-1028**  
 **EU-0024354-0000**

**UK authorisation holder:** Russell IPM,  
Unit 45, First Avenue, Deeside Industrial park,  
Deeside, Flintshire, UK CH5 2NU.

**EU authorisation holder:** Russell IPM Europe  
limited, Bond house 9/10 Bridge street,  
Lower Dublin 8, Dublin , Ireland.

**Active substance:**  
1433 – (9Z,12E)-tetradeca-9,  
12-dien-1-yl acetate 0,837%

**Type of Formulation:**  
VP-Vapour releasing product

## FIRST AID

**IF IN EYES:** If symptoms occur rinse with  
water. Remove contact lenses, if present  
and easy to do. Call a **POISON CENTRE**  
or a doctor.

**IF INHALED:** If symptoms occur call a  
**POISON CENTRE** or a doctor.

**IF ON SKIN:** Wash skin in water.  
If symptoms occur call a **POISON CENTRE**  
or a doctor.

**IF SWALLOWED:** If symptoms occur call  
a **POISON CENTRE** or a doctor.

**Target insects:**  
Indian meal moths (*Plodia interpunctella*)  
Mediterranean flour moths (*Ephesia*  
*Khueniella*)

**Where to use:**  
Cereal processing, bakeries, confectionary  
manufacturing facilities and dried food  
processing/storage facilities.

**Application:**  
One dispenser every 8m x 8m at 2m height  
(400-500 cubic meter). Change dispensers  
every three months.

**Target insects:**  
Indian meal moths (*Plodia interpunctella*)  
Mediterranean flour moths (*Ephesia*  
*Khueniella*)

**Where to use:**  
Cereal processing, bakeries, confectionary  
manufacturing facilities and dried food  
processing/storage facilities.

**Application:**  
One dispenser every 8m x 8m at 2m height  
(400-500 cubic meter). Change dispensers  
every three months.

Approved for the use in the presence of exposed (unpacked) food.

Approved for the use in the presence of exposed (unpacked) food.

**Dismate PE** acts by disrupting the mating  
between male and female moths. The  
populations of the pest insects are  
reduced as adult insects die and are not  
replaced by the next generation.  
**Dismate PE** is a vapourising dispenser in  
ready to use pouch. Keep the dispensers  
sealed in outer foil bag packaging until use.

**Store at temperatures not exceeding 4°C.**  
**Shelf life of up to 2 years.**

**Pack quantity: 5 or 100**

**DIRECTIONS FOR USE:**  
Plot out a layout for the approximate  
placement of the dispensers before deploying  
them. The dispensers should be placed one  
unit in every 400-500m<sup>3</sup>. Where obstacles or  
obstructions are present the dispensers can  
be placed to accommodate the layout and air  
dynamics of the factory.  
The dispensers should be positioned 2-3  
metres above floor and fastened securely  
using the supplied wall mounting bracket.  
Monitoring traps are recommended before,  
during and after treatment to assess

adult moth populations as part of a pest  
management program.  
Dispensers should be replaced every 3 months.  
Dispose of contents/container to national  
Regulations.  
This product has a slow mode of action and  
is intended for long-term use. The product  
doesn't have an immediate effect on the  
population of the food moths (it may take  
3-12 months to notice any significant  
reduction in moth population).  
Control of food moths is achieved over a period  
of time with continuous use of the product.

**Dismate PE** acts by disrupting the mating  
between male and female moths. The  
populations of the pest insects are  
reduced as adult insects die and are not  
replaced by the next generation.  
**Dismate PE** is a vapourising dispenser in  
ready to use pouch. Keep the dispensers  
sealed in outer foil bag packaging until use.

**Store at temperatures not exceeding 4°C.**  
**Shelf life of up to 2 years.**

**Pack quantity: 5 or 100**

**DIRECTIONS FOR USE:**  
Plot out a layout for the approximate  
placement of the dispensers before deploying  
them. The dispensers should be placed one  
unit in every 400-500m<sup>3</sup>. Where obstacles or  
obstructions are present the dispensers can  
be placed to accommodate the layout and air  
dynamics of the factory.  
The dispensers should be positioned 2-3  
metres above floor and fastened securely  
using the supplied wall mounting bracket.  
Monitoring traps are recommended before,  
during and after treatment to assess

adult moth populations as part of a pest  
management program.  
Dispensers should be replaced every 3 months.  
Dispose of contents/container to national  
Regulations.  
This product has a slow mode of action and  
is intended for long-term use. The product  
doesn't have an immediate effect on the  
population of the food moths (it may take  
3-12 months to notice any significant  
reduction in moth population).  
Control of food moths is achieved over a period  
of time with continuous use of the product.